

§ 341.90

Children under 6 years of age: consult a doctor.

(vi) *For products containing propylhexedrine identified in § 341.20(b)(9) when used in an inhalant dosage form.* The product delivers in each 800 milliliters of air 0.40 to 0.50 milligrams of propylhexedrine. Adults and children 6 to under 12 years of age (with adult supervision): 2 inhalations in each nostril not more often than every 2 hours. Children under 6 years of age: consult a doctor.

(vii) *For products containing xylometazoline hydrochloride identified in § 341.20(b)(10)—(A) Nasal drops or sprays—(1) For a 0.1-percent aqueous solution.* Adults and children 12 years of age and over: 2 or 3 drops or sprays in each nostril not more often than every 8 to 10 hours. Do not give to children under 12 years of age unless directed by a doctor.

(2) *A 0.05-percent aqueous solution in a container having either a calibrated dropper or a metered-dose spray that delivers no more than 0.054 milligrams of xylometazoline per three drops or three sprays.* Children 6 to under 12 years of age (with adult supervision): 2 or 3 drops or sprays in each nostril not more often than every 8 to 10 hours. Children 2 to under 6 years of age (with adult supervision): 2 or 3 drops or sprays in each nostril not more often than every 8 to 10 hours. Use only recommended amount. Do not exceed 3 doses in any 24-hour period. [previous two sentences in boldface type] Children under 2 years of age: consult a doctor.

(B) *Nasal jelly—(1) For a 0.1-percent water-based jelly.* Adults and children 12 years of age and over: place a small amount in each nostril and inhale well back into the nasal passages. Use not more often than every 8 to 10 hours. Do not give to children under 12 years of age unless directed by a doctor.

(2) *For a 0.05-percent water-based jelly.* Children 6 to under 12 years of age (with adult supervision): place a small amount in each nostril and inhale well back into the nasal passages. Use not more often than every 8 to 10 hours. Children under 6 years of age: consult a doctor.

(viii) *Other required statements—For products containing levmetamfetamine or*

21 CFR Ch. I (4–1–02 Edition)

propylhexedrine identified in § 341.20(b)(1) or (b)(9) when used in an inhalant dosage form. (A) “This inhaler is effective for a minimum of 3 months after first use.”

(B) “Keep inhaler tightly closed.”

[59 FR 43409, Aug. 23, 1994, as amended at 63 FR 40650, July 30, 1998; 64 FR 13295, Mar. 17, 1999; 65 FR 8, Jan. 3, 2000]

§ 341.90 Professional labeling.

The labeling of the product provided to health professionals (but not to the general public) may contain the following additional dosage information for products containing the active ingredients identified below:

(a) *For products containing ephedrine, ephedrine hydrochloride, ephedrine sulfate, or racephedrine hydrochloride identified in § 341.16 (a), (b), (c), and (f).* Children 6 to under 12 years of age: oral dosage is 6.25 to 12.5 milligrams every 4 hours, not to exceed 75 milligrams in 24 hours. Children 2 to under 6 years of age: oral dosage is 0.3 to 0.5 milligram per kilogram of body weight every 4 hours, not to exceed 2 milligrams per kilogram of body weight in 24 hours.

(b) *For products containing chlorphedianol hydrochloride identified in § 341.14(a)(1).* Children 2 to under 6 years of age: oral dosage is 12.5 milligrams every 6 to 8 hours, not to exceed 50 milligrams in 24 hours.

(c) *For products containing codeine ingredients identified in § 341.14(a)(2).* (1) Children 2 to under 6 years of age: Oral dosage is 1 milligram per kilogram body weight per day administered in four equal divided doses. The average body weight for each age may also be used to determine dosage as follows: For children 2 years of age (average body weight, 12 kilograms), the oral dosage is 3 milligrams every 4 to 6 hours, not to exceed 12 milligrams in 24 hours; for children 3 years of age (average body weight, 14 kilograms), the oral dosage is 3.5 milligrams every 4 to 6 hours, not to exceed 14 milligrams in 24 hours; for children 4 years of age (average body weight, 16 kilograms), the oral dosage is 4 milligrams every 4 to 6 hours, not to exceed 16 milligrams in 24 hours; for children 5 years of age (average body weight, 18 kilograms), the oral dosage is 4.5 milligrams every 4 to 6 hours, not to exceed 18 milligrams in

24 hours. The manufacturer must relate these dosages for its specific product dosages for its specific product to the use of the calibrated measuring device discussed in paragraph (c)(3) of this section. If age is used to determine the dose, the directions must include instructions to reduce the dose for low-weight children.

(2) Parents should be instructed to obtain and use a calibrated measuring device for administering the drug to the child, to use extreme care in measuring the dosage, and not exceed the recommended daily dosage.

(3) A dispensing device (such as a dropper calibrated for age or weight) should be dispensed along with the product when it is intended for use in children 2 to under 6 years of age to prevent possible overdose due to improper measuring of the dose.

(4) Codeine is not recommended for use in children under 2 years of age. Children under 2 years may be more susceptible to the respiratory depressant effects of codeine, including respiratory arrest, coma, and death.

(d) *The following labeling indication may be used for products containing guaifenesin identified in §341.18 when used as a single ingredient product.* “Helps loosen phlegm and thin bronchial secretions in patients with stable chronic bronchitis.”

(e) *For products containing brompheniramine maleate identified in §341.12(a).* Children 2 to under 6 years of age: oral dosage is 1 milligram every 4 to 6 hours, not to exceed 6 milligrams in 24 hours.

(f) *For products containing chlorcyclizine hydrochloride identified in §341.12(b).* Children 6 to under 12 years of age: oral dosage is 12.5 milligrams every 6 to 8 hours, not to exceed 37.5 milligrams in 24 hours. Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 6 to 8 hours, not to exceed 18.75 milligrams in 24 hours.

(g) *For products containing chlorpheniramine maleate identified in §341.12(c).* Children 2 to under 6 years of age: oral dosage is 1 milligram every 4 to 6 hours, not to exceed 6 milligrams in 24 hours.

(h) *For products containing dexbrompheniramine maleate identified in §341.12(d).* Children 2 to under 6 years

of age: oral dosage is 0.5 milligram every 4 to 6 hours, not to exceed 3 milligrams in 24 hours.

(i) *For products containing dexchlorpheniramine maleate identified in §341.12(e).* Children 2 to under 6 years of age: oral dosage is 0.5 milligram every 4 to 6 hours, not to exceed 3 milligrams in 24 hours.

(j) *For products containing diphenhydramine citrate identified in §341.12(f).* Children 2 to under 6 years of age: oral dosage is 9.5 milligrams every 4 to 6 hours, not to exceed 57 milligrams in 24 hours.

(k) *For products containing diphenhydramine hydrochloride identified in §341.12(g).* Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 mg in 24 hours.

(l) *For products containing doxylamine succinate identified in §341.12(h).* Children 2 to under 6 years of age: oral dosage is 1.9 to 3.125 milligrams every 4 to 6 hours, not to exceed 18.75 milligrams in 24 hours.

(m) *For products containing phenindamine tartrate identified in §341.12(i).* Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 milligrams in 24 hours.

(n) *For products containing pheniramine maleate identified in §341.12(j).* Children 2 to under 6 years of age: oral dosage is 3.125 to 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 milligrams in 24 hours.

(o) *For products containing pyrilamine maleate identified in §341.12(k).* Children 2 to under 6 years of age: oral dosage is 6.25 to 12.5 milligrams every 6 to 8 hours, not to exceed 50 milligrams in 24 hours.

(p) *For products containing thonzylamine hydrochloride identified in §341.12(l).* Children 2 to under 6 years of age: oral dosage is 12.5 to 25 milligrams every 4 to 6 hours, not to exceed 150 milligrams in 24 hours.

(q) *For products containing triprolidine hydrochloride identified in §341.12(m).* Children 4 to under 6 years of age: oral dosage is 0.938 milligram every 4 to 6 hours, not to exceed 3.744 milligrams in 24 hours. Children 2 to under 4 years of age: oral dosage is 0.625 milligram every 4 to 6 hours, not to exceed 2.5

milligrams in 24 hours. Infants 4 months to under 2 years of age: oral dosage is 0.313 milligram every 4 to 6 hours, not to exceed 1.252 milligrams in 24 hours.

(r) *For products containing diphenhydramine citrate identified in § 341.14(a)(5).* Children 2 to under 6 years of age: oral dosage is 9.5 milligrams every 4 hours, not to exceed 57 milligrams in 24 hours.

(s) *For products containing diphenhydramine hydrochloride identified in § 341.14(a)(6).* Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 hours, not to exceed 37.5 milligrams in 24 hours.

[51 FR 35339, Oct. 2, 1986, as amended at 52 FR 30057, Aug. 12, 1987; 54 FR 8509, Feb. 28, 1989; 57 FR 58376, Dec. 9, 1992; 59 FR 4218, Jan. 28, 1994; 59 FR 29174, June 3, 1994; 59 FR 36051, July 15, 1994]

PART 343—INTERNAL ANALGESIC, ANTIPYRETIC, AND ANTIRHEUMATIC DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

Sec.

343.1 Scope.

343.3 Definitions.

Subpart B—Active Ingredients

343.10 [Reserved]

343.12 Cardiovascular active ingredients.

343.13 Rheumatologic active ingredients.

343.20 [Reserved]

343.22 Permitted combinations of active ingredients for cardiovascular-rheumatologic use.

Subpart C—Labeling

343.50 [Reserved]

343.60 [Reserved]

343.80 Professional labeling.

Subpart D—Testing Procedures

343.90 Dissolution and drug release testing.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 63 FR 56814, Oct. 23, 1998, unless otherwise noted.

Subpart A—General Provisions

§ 343.1 Scope.

(a) An over-the-counter analgesic-antipyretic drug product in a form suitable for oral administration is generally recognized as safe and effective and is not misbranded if it meets each of the conditions in this part in addition to each of the general conditions established in § 330.1 of this chapter.

(b) References in this part to regulatory sections of the Code of Federal Regulations are to chapter I of title 21 unless otherwise noted.

§ 343.3 Definitions.

As used in this part:

Analgesic—antipyretic drug. An agent used to alleviate pain and to reduce fever.

Cardiovascular drug. An agent used to prevent ischemic events.

Rheumatologic drug. An agent used for the treatment of rheumatologic disorders.

Subpart B—Active Ingredients

§ 343.10 [Reserved]

§ 343.12 Cardiovascular active ingredients.

(a) Aspirin.

(b) Buffered aspirin. Aspirin identified in paragraph (a) of this section may be buffered with any antacid ingredient(s) identified in § 331.11 of this chapter provided that the finished product contains at least 1.9 milliequivalents of acid-neutralizing capacity per 325 milligrams of aspirin as measured by the procedure provided in the United States Pharmacopeia 23/National Formulary 18.

§ 343.13 Rheumatologic active ingredients.

(a) Aspirin.

(b) Buffered aspirin. Aspirin identified in paragraph (a) of this section may be buffered with any antacid ingredient(s) identified in § 331.11 of this chapter provided that the finished product contains at least 1.9 milliequivalents of acid-neutralizing capacity per 325 milligrams of aspirin as measured by the procedure provided in